AMENDMENTS TO THE CLAIMS

Claim 1 (Original). A compound of formula (I):



(I)

wherein:

m and n are independently 1 to 4;

R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl, haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;

R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl, carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;

R^3 is aryl or aryloxy each independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^8$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

or R^3 is aralkyl or aralkoxy, wherein the alkyl radical in the aralkyl or aralkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^9-N(R^7)C(O)OR^8$), and wherein the aryl radical in the aralkyl or aralkoxy substituent is independently optionally substituted by one or more

substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;

as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture; or a pharmaceutically acceptable salt thereof.

Claim 2 (Original). The compound of Claim 1 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aryl optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^8-N(R^7)C(O)OR^9$;

each R^4 is is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino,

hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

R^5 is hydrogen;

R^6 is hydrogen or alkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

R^9 is hydrogen, alkyl, aralkyl or haloalkyl.

Claim 3 (Original). The compound of Claim 2 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aryl optionally substituted by one or more substituents selected from the group consisting of carboxy or alkoxycarbonyl;

each R^4 is is independently selected from the group consisting of hydrogen, alkyl, halo, or haloalkyl;

R^5 is hydrogen; and

R^6 is hydrogen.

Claim 4 (Presently Amended). The compound ~~of Claim 4, namely,~~ 2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-carboxy)phenylquinoline ~~in trifluoroacetic acid~~, according to Claim 3.

Claim 5 (Original). The compound of Claim 1 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aryloxy optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

R^5 is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

R^9 is hydrogen, alkyl, aralkyl or haloalkyl.

Claim 6 (Original). The compound of Claim 5 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aryloxy optionally substituted by one or more substituents selected from the group consisting of alkyl, tetrazolyl, $-R^8-C(O)OR^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is is independently selected from the group consisting of hydrogen, alkyl, halo, or haloalkyl;

R^5 is hydrogen;

R^6 is hydrogen;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl; and

each R^8 is a bond or a straight or branched alkylene chain.

Claim 7 (Presently Amended). The compound of Claim 6 selected from the group consisting of the following:

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-carboxy)phenoxyquinoline ~~in-2,2,2-trifluoro-1,1-ethanediol~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(2-carboxy)phenoxyquinoline ~~in 2,2,2-trifluoro-1,1-ethanediol~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(2-amino-5-carboxy)phenoxyquinoline ~~in 2,2,2-trifluoro-1,1-ethanediol~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(4-carboxy)phenoxyquinoline ~~in 2,2,2-trifluoro-1,1-ethanediol~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-carboxymethyl)phenoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-(1-amino-1-carboxy)methyl)phenoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-(2-amino-2-carboxy)ethyl)phenoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(2-methyl-5-carboxy)phenoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(5-carboxy-2-diethylaminomethyl)phenoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-tetrazol-5-yl)phenoxyquinoline ~~in 2,2,2-trifluoro-1,1-ethanediol~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(3-trifluoromethylsulfonylamino)phenoxyquinoline ~~in trifluoroacetic acid~~; and

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-4-(3-carboxy)phenoxyquinoline ~~in trifluoroacetic acid~~.

Claim 8 (Original). The compound of Claim 1 wherein

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aralkyl wherein the alkyl radical in the aralkyl substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^9-N(R^7)C(O)OR^9$), and wherein the aryl radical in the aralkyl substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^9-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_z-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

R^5 is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

R^9 is hydrogen, alkyl, aralkyl or haloalkyl.

Claim 9 (Original). The compound of Claim 1 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aralkoxy wherein the alkyl radical in the aralkyl substituent is not optionally substituted and wherein the aryl radical in the aralkoxy substituent is optionally substituted by

one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

R^5 is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

R^9 is hydrogen, alkyl, aralkyl or haloalkyl.

Claim 10 (Original). The compound of Claim 9 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aralkoxy wherein the aryl radical in the aralkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, and $-R^8-N(R^7)_2$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, halo, or haloalkyl;

R^5 is hydrogen;

R^6 is hydrogen;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl; and

each R^8 is a bond or a straight or branched alkylene chain.

Claim 11 (Presently Amended). The compound ~~of Claim 10~~ selected from the group consisting of the following:

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-4-benzyloxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-benzyloxycarbonylpropyl]aminocarbonyl-4-benzyloxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-benzyloxyquinoline;

2-[1-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-benzyloxycarbonylpropyl]aminocarbonyl-4-benzyloxy-8-methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-benzyloxy-8-methoxyquinoline;

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-4-(4-methoxycarbonyl)benzyloxyquinoline;

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-4-(4-carboxy)benzyloxyquinoline;

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-4-(3-methoxycarbonyl)benzyloxyquinoline;

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-4-(3-carboxy)benzyloxyquinoline;

2-[1*S*-(4-(3-methylphenyl)piperazin-1-yl)carbonyl-3-(1,1-dimethylethoxycarbonyl)propyl]aminocarbonyl-4-benzyloxyquinoline; and

2-[1*S*-(4-(3-methylphenyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-benzyloxyquinoline.

Claim 12 (Original). The compound of Claim 1 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aralkoxy wherein the alkyl radical in the aralkoxy substituent is substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^6-N(R^7)C(O)R^7$, and $-R^9-N(R^7)C(O)OR^9$), and wherein the aryl radical in the aralkoxy substituent is

optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

R^5 is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

R^9 is hydrogen, alkyl, aralkyl or haloalkyl.

Claim 13 (Original). The compound of Claim 12 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is aralkoxy wherein the alkyl radical in the aralkoxy substituent is substituted by $-R^8-C(O)OR^7$, and wherein the aryl radical in the aralkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo and $-R^8-OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, halo, haloalkyl, amino, monoalkylamino, or dialkylamino;

R^5 is hydrogen;

R^6 is hydrogen;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl; and

each R^8 is a bond or a straight or branched alkylene chain.

Claim 14 (Presently Amended). The compound of Claim 13 selected from the group consisting of the following:

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-(1,1-dimethylethoxycarbonyl)propyl]aminocarbonyl-4-(1-phenyl-1-methoxycarbonyl)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(1-phenyl-1-methoxycarbonyl)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-chloro-4-(1-carboxy-1-phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-naphth-1-yl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-chloro-8-fluoro-4-(1-methoxycarbonyl-1-phenyl)methoxyquinoline ~~in acetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-chloro-8-fluoro-4-(1-carboxy-1-phenyl)methoxyquinoline ~~in acetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-6-fluoro-4-(1-carboxy-1-(2-fluoro)phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-4-(1-ethoxycarbonyl-1-phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[(4-(ethoxycarbonyl)piperazin-1-yl)carbonylmethyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-6-fluoro-4-(1-carboxy-1-(4-chloro)phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-6-fluoro-4-(1-carboxy-1-(3-methoxy)phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6,8-difluoro-4-(1-carboxy-1-phenyl)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-dimethylamino-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-chloro-6-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-7-methyl-6-chloro-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(1,1-dimethylethoxycarbonyl)piperazin-1-yl)carbonyl-3-methoxycarbonylpropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(1,1-dimethylethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(methoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(1,1-dimethylethylaminocarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(furan-2-ylcarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(3-methylphenyl)piperazin-1-yl)carbonyl-3-(1,1-dimethylethoxycarbonyl)propyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline;

2-[1*S*-(4-(3-methylphenyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~;

2-[1*S*-(4-(phenyl)piperazin-1-yl)carbonyl-3-(1,1-dimethylethoxycarbonyl)propyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline; and

2-[1*S*-(4-(phenyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-6-fluoro-7-methyl-4-(1-phenyl-1-carboxy)methoxyquinoline ~~in trifluoroacetic acid~~.

Claim 15 (Original). A pharmaceutical composition useful in treating a mammal having a disease-state characterized by thrombotic activity, which composition comprises a pharmaceutically acceptable excipient and a compound of formula (I):



(I)

wherein:

m and n are independently 1 to 4;

R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl, haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;

R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl, carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;

R^3 is aryl or aryloxy each independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

or R^3 is aralkyl or aralkoxy, wherein the alkyl radical in the aralkyl or aralkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^9-N(R^7)C(O)OR^9$), and wherein the aryl radical in the aralkyl or aralkoxy substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$,

$-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;

as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture; or a pharmaceutically acceptable salt thereof.

Claim 16 (Original). A method of treating a disease-state characterized by thrombotic activity, which method comprises administering to a mammal having a disease-state characterized by thrombotic activity a therapeutically effective amount of a compound of formula (I):



(I)

wherein:

m and n are independently 1 to 4;

R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl,

haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;

R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl, carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;

R^3 is aryl or aryloxy each independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

or R^3 is aralkyl or aralkoxy, wherein the alkyl radical in the aralkyl or aralkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^8-N(R^7)C(O)OR^9$), and wherein the aryl radical in the aralkyl or aralkoxy substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;
each R^8 is a bond or a straight or branched alkylene chain; and
each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;
as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture;
or a pharmaceutically acceptable salt thereof.

Claim 17 (Presently Amended). A compound of formula (II):



(II)

wherein:
m and n are independently 1 to 4;
R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl, haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;
R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl, carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;
R^3 is heteroaryl optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

~~or R^3 is heteroarylalkoxy, wherein the alkoxy radical in the heteroarylalkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, R^8-OR^7, R^8-$C(O)OR^7$, R^8-$C(O)N(R^7)_2$, R^8-$C(O)R^7$, R^8-$N(R^7)_2$, R^8-$N(R^7)C(O)R^7$, and R^9-$N(R^7)C(O)OR^9$), and wherein the heteroaryl radical in the heteroarylalkoxy substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, R^8-OR^7, R^8-$C(O)OR^7$, R^8-$C(O)N(R^7)_2$, R^8-$C(O)R^7$, R^8-$N(R^7)_2$, R^8-$N(R^7)C(O)R^7$, R^8-$N(R^7)C(O)OR^9$, R^8-$N(R^7)$-$S(O)_2$-R^7, and R^8-$C[N(R^7)_2]$-$C(O)OR^7$;~~

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;

as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture;

or a pharmaceutically acceptable salt thereof.

Claim 18 (Original). The compound of Claim 17 wherein:

m is 1;

n is 1 or 2;

R^1 is hydrogen, aryl, aralkyl, or alkoxycarbonyl;

R^2 is hydrogen, carboxyalkyl, alkoxycarbonylalkyl or aralkoxycarbonylalkyl;

R^3 is heteroaryl optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, -R^8-OR^7, -R^8-$C(O)OR^7$, -R^8-$C(O)N(R^7)_2$, and -R^8-$N(R^7)_2$;

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, halo, haloalkyl, amino, monoalkylamino, or dialkylamino;

R^5 is hydrogen;

R^6 is hydrogen;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl; and

each R^8 is a bond or a straight or branched alkylene chain.

Claim 19 (Presently Amended). The compound ~~of Claim 18, namely,~~ 2-[1*S*-(4-(ethoxycarbonyl)piperazin-1-yl)carbonyl-3-carboxypropyl]aminocarbonyl-4-(1,2,3,4-tetrahydroisoquinolin-2-yl)quinoline ~~in trifluoroacetic acid~~, **according to Claim 18**.

Claim 20 (Canceled).

Claim 21 (Canceled).

Claim 22 (Presently Amended). A pharmaceutical composition useful in treating a mammal having a disease-state characterized by thrombotic activity, which composition comprises a pharmaceutically acceptable excipient and a compound of formula (II):



(II)

wherein:

m and n are independently 1 to 4;

R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl, haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;

R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl,

carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;

R^3 is heteroaryl optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

~~or R^3 is heteroarylalkoxy, wherein the alkoxy radical in the heteroarylalkoxy substituent is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^8-N(R^7)C(O)OR^9$), and wherein the heteroaryl radical in the heteroarylalkoxy substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;~~

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;

as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture;

or a pharmaceutically acceptable salt thereof.

Claim 23 (Presently Amended). A method of treating a disease-state characterized by thrombotic activity, which method comprises administering to a mammal having a disease-state characterized by thrombotic activity a therapeutically effective amount of a compound of formula (II):



(II)

wherein:

m and n are independently 1 to 4;

R^1 is hydrogen, alkyl, carboxyalkyl, aryl, aralkyl, alkylcarbonyl, aryloxyalkylcarbonyl, carboxyalkylcarbonyl, alkoxycarbonylalkylcarbonyl, alkoxycarbonylalkyl, alkoxycarbonyl, arylcarbonyl, aryloxycarbonyl, aralkoxycarbonyl, cycloalkylcarbonyl, haloalkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkoxycarbonylaminocarbonyl, or heterocyclylcarbonyl;

R^2 is hydrogen, alkyl, aryl, aralkyl, alkylsulfonylalkyl, aralkoxyalkyl, hydroxyalkyl, aminoalkyl, haloalkylsulfonylaminoalkyl, carboxyalkylthioalkyl, alkoxycarbonylalkylthioalkyl, carboxyalkyl, (carboxy)(hydroxy)alkyl, carboxyalkoxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, carboxyalkoxycarbonylalkyl, alkoxycarbonylalkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxycarbonylaminoalkyl, alkoxycarbonylalkylaminocarbonylalkyl, carboxyalkylaminocarbonylalkyl, (alkoxycarbonylalkyl)(alkyl)aminocarbonylalkyl, (carboxyalkyl)(alkyl)aminocarbonylalkyl, or heterocyclylalkyl;

R^3 is heteroaryl optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;

~~or R^3 is heteroarylalkoxy, wherein the alkoxy radical in the heteroarylalkoxy substituent is optionally substituted by one or more substituents selected from~~

~~the group consisting of halo, cyano, nitro, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, and $-R^9-N(R^7)C(O)OR^9$, and wherein the heteroaryl radical in the heteroarylalkoxy substituent is independently optionally substituted by one or more substituents selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, tetrazolyl, $-R^8-OR^7$, $-R^8-C(O)OR^7$, $-R^8-C(O)N(R^7)_2$, $-R^8-C(O)R^7$, $-R^8-N(R^7)_2$, $-R^8-N(R^7)C(O)R^7$, $-R^8-N(R^7)C(O)OR^9$, $-R^8-N(R^7)-S(O)_2-R^7$, and $-R^8-C[N(R^7)_2]-C(O)OR^7$;~~

each R^4 is independently selected from the group consisting of hydrogen, alkyl, alkoxy, aralkoxy, halo, haloalkyl, haloalkoxy, hydroxy, cyano, alkylthio, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyl, nitro, amino, monoalkylamino, dialkylamino, carboxyalkylamino, alkylcarbonylamino, di(alkylcarbonyl)amino, hydroxyalkyl, dialkylaminoalkyl, carboxyalkoxy, alkoxycarbonylalkoxy, dialkylaminoalkoxy, and heterocyclylalkoxy;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, aralkyl, carboxy, alkoxycarbonyl, aralkoxycarbonyl, carboxyalkyl, and alkoxycarbonylalkyl;

R^6 is hydrogen, alkyl, carboxyalkyl, or alkoxycarbonylalkyl;

each R^7 is hydrogen, alkyl, aryl, aralkyl, or haloalkyl;

each R^8 is a bond or a straight or branched alkylene chain; and

each R^9 is hydrogen, alkyl, aralkyl or haloalkyl;

as a single stereoisomer, a mixture of individual stereoisomers, or a racemic mixture; or a pharmaceutically acceptable salt thereof.